

**AIC Endocrinology #1****Fax: 833-808-0833****Patient Information**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Height: \_\_\_\_\_ Weight: \_\_\_\_\_ Patient Contact #: \_\_\_\_\_

**Medical Assessment**

Diagnosis:

- ☐ M80.0\_\_ Age related Osteoporosis with current fracture  
☐ M81.0 Age related Osteoporosis without current fracture  
☐ C61 Malignant Neoplasm of the Prostate  
☐ C50.\_\_\_\_ Breast Cancer  
☐ Other: \_\_\_\_\_

Failed Medications: \_\_\_\_\_

Has the patient tried and failed oral bisphosphonates? If so, which ones and when? \_\_\_\_\_

Dexa Scan T Score: \_\_\_\_\_ Type: \_\_\_\_\_ Date: \_\_\_\_\_

Fracture History: Site \_\_\_\_\_ Date: \_\_\_\_\_

Thyrogen Use: (Please mark)

- ☐ Use as an adjunctive treatment as pre-therapeutic stimulation for radioiodine ablation of thyroid tissue remnants in patients maintained on thyroid hormone suppression therapy who have undergone a near-total or total thyroidectomy for well-differentiated thyroid cancer without evidence of distant metastatic thyroid cancer.  
☐ Use as an adjunctive diagnostic tool for serum thyroglobulin (Tg) testing, with or without radioiodine imaging, in the follow-up of patients with well-differentiated thyroid cancer  
☐ Other: \_\_\_\_\_

**Pre-Medication Orders:**

- ☐ APAP ☐ 325mg ☐ 500mg ☐ 650mg - PO 30 minutes before infusion  
☐ Diphenhydramine ☐ 25mg ☐ 50mg – PO 30 minutes before infusion OR ☐ 25mg IVP ☐ 50mg IVP  
☐ Alternate Oral Antihistamines: ☐ Cetirizine 10mg ☐ Loratadine 10mg ☐ Fexofenadine 60mg or ☐ 180mg  
☐ Methylprednisolone ☐ 40mg IVP ☐ 125mg IVP or other \_\_\_\_\_ mg IVP  
☐ Famotidine ☐ 20mg PO ☐ 40mg PO ☐ 20mg IVP ☐ 40mg IVP  
☐ Ondansetron ☐ 4 mg IVP ☐ 4 mg PO

**Labwork:** \_\_\_\_\_ To be Drawn by: ☐ Infusion Clinic Referring Provider**Allergies:** \_\_\_\_\_**Medication****Thyrogen (thyrotropin alpha)**

Dosage / Frequency:

- ☐ Thyrogen 0.9mg IM on day 1 and 2.

**Evenity (romosozumab-aqqg)**

Dosage / Frequency:

- ☐ Evenity 210mg SQ monthly for 12 months.

**Maintenance Refills:** \_\_\_\_\_**Ibandronate Sodium (generic Boniva)**

Dosage / Frequency:

- ☐ Ibandronate sodium 3 mg IV every 3 months.

**Maintenance Refills:** \_\_\_\_\_**Physician Prescription Orders** Address: \_\_\_\_\_ Phone: \_\_\_\_\_

Fax: \_\_\_\_\_ Clinic Contact: \_\_\_\_\_ Physician Name: \_\_\_\_\_

NPI # \_\_\_\_\_ **Physician Signature:** \_\_\_\_\_ **Date:** \_\_\_\_\_