

**Patient Information**

Patient Name: _____ DOB: _____ Height: _____ Weight: _____ Patient Contact #: _____

Medical Assessment1. **Diagnosis:** ☐ M06.9 Rheumatoid Arthritis ☐ L40.50 Psoriatic Arthritis ☐ Ankylosing Spondylitis ☐ Other: _____
☐ M32.10 Systemic lupus erythematosus ☐ M1A.__0 Chronic gout ☐ M10.__ Idiopathic gout2. **Drug Allergies:** _____3. **Failed Medications:** ☐ Methotrexate Therapy Length: _____ Discontinuation Reason: _____
☐ _____ Therapy Length: _____ Discontinuation Reason: _____****Please include a demographics page, front and back of an insurance card-medical / RX insurance, any clinical notes with supporting diagnosis, lab-work, tests, and any other supporting documentation including past tried and / or failed therapies associated with diagnosis******Labwork:** _____ To be Drawn by: Infusion Clinic Referring Provider**Pre-Medication Orders**

- ☐ APAP ☐ 325mg ☐ 500mg ☐ 650mg - PO 30 minutes before infusion.
- ☐ Diphenhydramine ☐ 25mg ☐ 50mg – PO 30 minutes before infusion OR 25mg IVP ☐ 50mg IVP
- ☐ Alternate Oral Antihistamines: ☐ Cetirizine 10mg ☐ Loratadine 10mg ☐ Fexofenadine 60mg or ☐ 180mg
- ☐ Methylprednisolone ☐ 40mg IVP ☐ 125mg IVP or other _____ mg IVP
- ☐ Famotidine ☐ 20mg PO ☐ 40mg PO ☐ 20mg IVP ☐ 40mg IVP
- ☐ Ondansetron ☐ 4 mg IVP ☐ 4 mg PO

Medication**Infliximab (Remicade & Biosimilars)**☐ **Preferred - Infusion Clinic Preference (Remicade / Avsola / Inflectra / Renflexis) based on payer directives / availability**☐ **Specific Product:** _____

Dosage: 3 mg /kg OR _____ mg or _____ mg / kg

Frequency:

- ☐ Induction: at week 0, 2, and 6 weeks
- ☐ Maintenance: at week 14 and every 8 weeks
- ☐ Other orders: _____
- ☐ **Maintenance Refills:** ____

Orencia (abatacept)Dosage: ☐ 500mg ☐ 750mg ☐ 1000mg

Frequency:

- ☐ Induction: at week 0, 2 and, 4
- ☐ Maintenance: at week 8 and every 4 weeks thereafter
- ☐ **Maintenance Refills:** ____

Rituximab (Rituxan & Biosimilars)☐ **Preferred - Infusion Clinic Preference (Rituxan / Ruxience / Truxima) based on payer directives / availability**☐ **Specific Product:** _____

Dosage: 1000mg IV

Frequency:

- ☐ Infuse at week 0 and 2, once every 4 months (16 weeks)
- ☐ Infuse at week 0 and 2, once every 6 months (24 weeks)
- ☐ Other orders: _____
- ☐ **Maintenance Refills:** ____

**** Pre-medications of acetaminophen PO, diphenhydramine IVP, and methylprednisolone is recommended prior to infusion******Physician Prescription Orders** Address: _____ Phone: _____

Fax: _____ Clinic Contact: _____ Physician Name: _____

NPI # _____ **Physician Signature:** _____ **Date:** _____